

IROSPAN® 24/6 Iron / Folic Acid / Supplement Tablets and Succinic Acid Tablets provide a 30 day oral iron supplement therapy

Irospan Tablet Description:

Each light blue capsule-shaped film coated tablet has a pleasant sweet flavor with “WC-002” imprinted on one side:

Ingredients:	Dose	Daily Value (2000 Calorie Diet)
Vitamin C (Sodium Ascorbate)	100 mg	167%
Formical® (Calcium Formate)	155 mg	16%
B1 Thiamine HCL	5 mg	333%
B2 Riboflavin	5 mg	294%
B3 Niacin	25 mg	125%
B6 Pyridoxine HCL	30 mg	1500%
Folic Acid	1000 mcg	250%
B12 Cyanocobalamin	10 mcg	167%
B7 Biotin	300 mcg	100%
B5 d-Calcium Pantothenate	7 mg	70%
Iron (Ferrous bisGlycinate Chelate)	65 mg	361%
Iron (Polysaccharide Iron Complex)	65 mg	361%

Inactive ingredients: Microcrystalline Cellulose, Magnesium Stearate, Povidone, Polyvinyl Alcohol-Partially Hydrolyzed, PEG 3350, Titanium Dioxide, Talc, FD&C Blue #1, Saccharin, Carnauba Wax.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.

Succinic Acid Tablet Description:

Each white round film coated tablet has a pleasant sweet flavor with “WC” imprinted on one side:

Ingredients:	Dose	Daily Value (2000 Calorie Diet)
Vitamin C (as Sodium Ascorbate)	100 mg	167%
Calcium (Calcium Carbonate)	100 mg	10%
B1 Thiamine HCL	5 mg	333%
B2 Riboflavin	5 mg	294%
B3 Niacin	25 mg	125%
B6 Pyridoxine HCL	30 mg	1500%
Folic Acid	1000 mcg	250%
B12 Cyanocobalamin	10 mcg	167%
B7 Biotin	300 mcg	100%
B5 d-Calcium Pantothenate	7 mg	70%
Succinic Acid	150 mg	*

- DV not established

Inactive ingredients: Microcrystalline Cellulose, Magnesium Stearate, Povidone, Polyvinyl Alcohol-Partially Hydrolyzed, PEG 3350, Titanium Dioxide, Talc, Saccharin, Carnauba Wax.

Clinical Pharmacology:

Iron is an essential component in the formation of hemoglobin. Adequate amounts of iron are necessary for effective erythropoiesis. Iron also serves as a cofactor of several essential enzymes, including cytochromes that are involved in electron transport.

Folic acid is required for nucleoprotein synthesis and the maintenance of normal erythropoiesis. Folic acid is converted in the liver and plasma to its metabolically active form, tetrahydrofolic acid, by dihydrofolate reductase.

Vitamin B₁₂ is required for the maintenance of normal erythropoiesis, nucleoprotein and myelin synthesis, cell reproduction and normal growth. Intrinsic factor, a glycoprotein secreted by the gastric mucosa, is required for active absorption of Vitamin B₁₂ from the gastrointestinal tract.

Indications and Usage:

IROSPAN[®] 24/6 is indicated for the treatment of all anemias that are responsive to oral iron therapy. These include: hypochromic anemia associated with pregnancy, chronic and/or acute blood loss, metabolic disease, post-surgical convalescence, and dietary needs.

Contraindications:

This product is contraindicated in patients with a known hypersensitivity to any of the ingredients. Hemolytic anemia, hemochromatosis, and hemosiderosis are contraindications to iron therapy.

Warnings:

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where vitamin B₁₂ is deficient.

Precautions:

Folic acid in doses above 0.1mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. This product contains formate, which if consumed above the recommended level, could cause visual impairment and other health effects. Do not take more than the recommended amount. If you are pregnant, nursing, or taking any medications consult your doctor before use. Discontinue use and consult your doctor if any adverse reactions occur. Not intended for use by persons under the age of 18.

Adverse Reactions:

Adverse reactions with iron therapy may include constipation, diarrhea, nausea, vomiting, dark stools, and abdominal pain. Adverse reactions with iron therapy are usually transient. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

Dosage and Administration:

Usual adult dosage is 1 tablet daily or as directed by a physician. The IROSPAN[®] 24/6 carton contains a 30 day course of iron therapy that consists of 24 light blue tablets and 6 white tablets. Take 1 light blue tablet daily for 24 days, followed by 1 white tablet daily for 6 days. After the 30 tablets have been taken, a new course may be started if prescribed.

How Supplied:

IROSPAN[®] 24/6 tablets for oral administration are available in four child resistant blister cards containing 30 tablets (24 light blue tablets and 6 white tablets) (NDC 50967-126-30)

Store at controlled room temperature 15°-30°(59°-86°F) **RX only**

Distributed by: Women's Choice Pharmaceuticals • Gilbert, AZ 85233 • (877) 774-4949

Formical[®] is a registered trademark of Nephro-Tech 1, LLC
US Patent #6,528,542, #6,489,361, #6,160,016

IROSPAN[®] is a registered trademark of Women's Choice Pharmaceuticals, LLC